

SHEP 01-30
February 1, 2008

Supersymmetric Inflation, Matter and Dark Energy

S. F. KING

*Department of Physics and Astronomy,
University of Southampton,
Southampton, SO17 1BJ, U.K.*

In this talk, based on [1], we emphasise that intermediate scale supersymmetric inflation models are particularly attractive since inflation, baryogenesis and the relic abundance of cold dark matter (CDM) are all related by a set of parameters which also affect particle physics collider phenomena, neutrino masses and the strong CP problem. We also point out that the present day relic abundances of different forms of matter are (in principle) calculable from the supersymmetric inflation model together with a measurement of the CMB temperature and the Hubble constant. From these relic abundances one can deduce the amount of the present day dark energy (DE) density.

PRESENTED AT

COSMO-01

Rovaniemi, Finland,
August 29 – September 4, 2001

1 Introduction

The introductory parts of the talk included the motivation for supersymmetry and inflation, and ten challenges facing a supersymmetric inflationary model. This was followed by a detailed review of a particular intermediate scale supersymmetric inflation model, as an example of a model which attempts to meet all the challenges. These parts of my talk can be found in the recent review article [2] to which we refer those readers requiring more background information. Here we state the omega problem in supersymmetric inflation, give the motivation for intermediate scale inflation and introduce a particular model. We then show how the parameters of this model simultaneously control different phenomena in particle physics and cosmology. Finally we show how the matter relic densities may be calculated from such a model together with a measurement of the CMB temperature and the Hubble constant, and hence how the DE density may be deduced.

2 The Omega Problem in Supersymmetric Inflation

Within the framework of intermediate scale supersymmetric inflation one may expect on general grounds that our Universe contains sizeable relic abundances of baryons (from e.g. leptogenesis), axions (a) (from the Peccei-Quinn (PQ) solution to the strong CP problem), as well as weakly interacting massive particles (WIMPS). In R-parity conserving SUSY the WIMP is identified as the lightest supersymmetric particle (LSP), and this is often assumed to be the lightest neutralino of the minimal supersymmetric standard model (MSSM). We shall argue from this perspective that the LSP could equally well be a lighter stable singlet (singlino) identified with an axino or inflatino.

In general the ratio of the total density of the universe ρ_{tot} to critical density ρ_{crit} is given by Ω_{tot} where

$$\Omega_{tot} = \Omega_{\gamma} + \Omega_{matter} + \Omega_{DE} \quad (1)$$

and Ω_{γ} , Ω_{matter} , Ω_{DE} are the ratios of radiation density ρ_{γ} , matter density ρ_{matter} and DE density ρ_{DE} to critical density ρ_{crit} , and the radiation density is unimportant $\rho_{\gamma} \ll \rho_{matter}$. Note that the critical density is a function of time and in the present epoch $\rho_{crit} = 3M_P^2 H_0^2 = (3h^{1/2} \times 10^{-3} \text{ eV})^4 \sim (M_W^2/M_P)^4$ where M_W is the weak scale, M_P is the Planck scale, and $H_0 = 100h \text{ km.s}^{-1} \text{ Mpc}^{-1}$ is the present day Hubble constant, with $h = 0.7 \pm 10\%$. From observation $\Omega_{DE} \sim 2/3$ while $\Omega_{matter} \sim 1/3$ and Ω_{tot} is very close to unity. Inflation predicts $\Omega_{tot} = 1$, for all times after inflation. The matter contributions consist of (at least)

$$\Omega_{matter} = \Omega_b + \Omega_{\nu} + \Omega_{LSP} + \Omega_a \quad (2)$$

The most recent data is consistent with nucleosynthesis estimates of $\Omega_b \sim 0.04$, where the baryons (b) in the universe are mainly to be found in dark objects. The determined value of $\Omega_{CDM} \sim 0.3$ contains unknown relative contributions from Ω_{LSP} and Ω_a . Super-Kamiokande sets a lower limit on neutrino masses $\sum_i m_{\nu_i} \geq 0.05 \text{ eV}$ which corresponds

to $\Omega_\nu \geq 0.003$. In hierarchical neutrino mass models the lower bound is saturated and interestingly the neutrino density is then comparable to the visible baryon density $\Omega_{stars} \sim 0.005$.

3 An Example

3.1 Why Intermediate Scale Inflation?

We now wish to consider a specific example of a model which addresses the particle physics issues mentioned in the abstract, in order to illustrate many of the general features that we have discussed above. The brand of inflation most closely related to particle physics seems to be hybrid inflation which may occur at a scale well below the Planck scale, and hence be in the realm of particle physics. The next question is what is the relevant scale at which hybrid inflation takes place? One obvious possibility is to associate the scale of inflation with some grand unified theory (GUT) symmetry breaking scale, as originally conceived by Guth. However it is somewhat ironic that hybrid inflation at the GUT scale faces the magnetic monopole problem, which was precisely one of the original motivations for considering inflation in the first place! Although in certain cases this problem may be resolved, there are typically further symmetry breaking scales below the GUT scale at which discrete symmetries are broken, leading to problems with cosmological domain walls.

Intermediate scale hybrid inflation immediately solves both the magnetic monopole problem and the domain wall problem. The idea is simply that there is a period of hybrid inflation occurring below the GUT scale at the PQ symmetry breaking scale itself, in which the inflaton carries PQ charge and so the choice of domain is fixed during inflation. The universe therefore inflates inside a particular domain, and the magnetic monopole relics produced by the GUT scale symmetry breaking are inflated away. This provides a powerful motivation for intermediate scale inflation.

3.2 Intermediate Scale Supersymmetric Inflation Model

The model we consider is a variant of the NMSSM. This model has a SUGRA foundation and leptogenesis and reheating has been studied and preheating has been demonstrated not to lead to over-production of either axions or gravitinos. The model provides a solution to the strong CP problem and the μ problem, with inflation directly solving the monopole and domain wall problems at the inflation scale. It is therefore a well motivated, successful model that has been well studied and does not appear to suffer from any embarrassing problems, and is therefore a suitable laboratory for our discussion here. This variant of the NMSSM has the following superpotential terms involving the standard Higgs doublets H_u, H_d and two gauge singlet fields ϕ (inflaton) and N ,

$$W = \lambda N H_u H_d - k \phi N^2 \quad (3)$$

where λ, k are dimensionless coupling constants. Notice that the standard NMSSM is recovered if we replace the inflaton ϕ by N . However this leads to the familiar domain wall problems arising from the discrete Z_3 symmetry. In this new variant, the Z_3 becomes a global $U(1)_{PQ}$ symmetry that is commonly invoked to solve the strong CP problem. This symmetry is broken in the true vacuum by intermediate scale ϕ and N VEVs, where the axion is the pseudo-Goldstone boson from the spontaneous symmetry breaking and constrains the size of the VEVs. With such large VEVs this model should be regarded as giving an intermediate scale solution to the μ problem, and as such will have collider signatures.

We can make the ϕ -field real by a choice of the (approximately) massless axion field. We will now regard ϕ and N to be the real components of the complex singlets in what follows. When we include soft SUSY breaking mass terms, trilinear terms $A_k k \phi N^2 + h.c.$ (for real A_k) and neglect the $\lambda N H_u H_d$ superpotential term, we have the following potential:

$$V = V_0 + k^2 N^4 + \frac{1}{2} m^2(\phi) N^2 + \frac{1}{2} m_\phi^2 \phi^2 \quad (4)$$

where $m^2(\phi) = m_N^2 + 4k^2 \phi^2 - 2k A_k \phi$. We can identify the various elements of the potential: V_0 arises from some other sector of the theory, SUGRA for example, and dominates the potential; the soft SUSY breaking parameters A_k and m_N are generated through some gravity-mediated mechanism with a generic value of $O(TeV)$.

More details about how hybrid inflation may be implemented in this model may be found in [2]. We only remark here that it is non-trivial that a set of parameters exists that is consistent with axion and SUSY physics and allows the correct COBE perturbations to be achieved by radiative corrections. Without SUSY one would be free to add soft scalar masses at will, but with SUSY one must rely on the theory which either generates soft masses of order a TeV, or sets them equal to zero as in no-scale SUGRA, in which case the radiative corrections, which are under control in the case of SUSY predict the relevant value of the soft parameters, without any further adjustable parameters. Thus SUSY is playing a crucial role in the model which is why we refer to it as a Supersymmetric Inflation Model.

3.3 The Cosmological Constant Problem

Notice that the SUGRA-derived potential contribution V_0 exactly cancels with the other terms (by tuning) to provide agreement with the observed small cosmological constant. Thus we assume that at the global minimum $V(\langle\phi\rangle, \langle N\rangle) = 0$ which implies that $V_0 = k^2 \langle N \rangle^4$. The height of the potential during inflation is therefore $V_0^{1/4} = k^{1/2} \langle N \rangle \sim 10^8 GeV$. Since the approach has a consistent way to set the large cosmological constant to zero, the absence of a real solution to this problem may not be an obstacle to the implications of the approach.

	V_0	k	λ	$\mathcal{L}_{soft}$	$\mathcal{L}_{Yuk}$
Inflation and COBE	✓	✓	-	✓	-
MSSM μ parameter	-	-	✓	✓	-
Fermion Masses, Mixings	-	-	-	-	✓
SUSY collider physics	-	✓	✓	✓	✓
Strong CP, axion abundance (Ω_a)	-	✓	✓	✓	-
Leptogenesis (Ω_b)	-	✓	✓	✓	✓
LSP abundance (Ω_{LSP})	-	✓	✓	✓	-

Table 1: This table illustrates the fact that a particular parameter of the model (columns) simultaneously controls several different aspects of particle physics and cosmology (rows) which are thereby related. $\mathcal{L}_{soft}$ contains A_k, m_N^2 and the other soft parameters, $\mathcal{L}_{Yuk}$ contains the Yukawa coupling constants controlling all fermion masses and mixing angles.

3.4 Parameter Counting and Singlino Mixing

A relevant parameter count at this stage reveals two superpotential effective parameters (λ and k), the two soft SUSY breaking parameters (A_k and m_N), plus the constant energy density V_0 . From these five parameters we have inflated the universe with the correct COBE perturbations, provided a μ term of the correct order of magnitude and solved the strong CP problem. They also govern the phenomenology of the singlet Higgs and Higgsino components of ϕ and N which may weakly mix with the MSSM superfields H_u, H_d .

The LSP will be the lightest eigenvalue of the full “ino” matrix, extended in the usual way to include gaugino-higgsino mixing. Clearly if $k < \lambda/2$ then a singlino will be the LSP. In our case the singlino may be regarded as a linear combination of axino and inflatino.

As usual in models based on an intermediate scale solution to the μ problem the coupling of the singlino to the neutralinos means that $\tilde{S}$ nearly decouples. However the conservation of R-parity means that eventually the lightest neutralino produced in colliders must decay into the singlino, and all the collider signatures discussed in may apply. In the case that the lightest neutralino leaves the detector before it decays into the singlino, there will be no unconventional collider signature. In this case the knowledge concerning a lighter singlino will come from cosmology since the LSP relic density gets diluted by the ratio of the singlino to lightest neutralino masses, and direct dark matter searches will not see anything since the singlino LSP will not scatter off nuclei.

One of the main things emphasized in [1] is the connection between the calculation of relic densities and the other physics, via their common parameters. This is summarised in Table 1 for the particular model we have been discussing. The same parameters that control the ino mass matrix will also be involved in reheating of the universe after inflation, and giving the relic densities of LSP and in leptogenesis as we discussed in [1]. Different models may have different mechanisms to solve some of the problems, different reheating and preheating, and so on, but will still lead to a version of Table 1.

4 How To Calculate the Size of the Dark Energy Density in Supersymmetric Inflation

Is there anything that we can say about DE at the current time from the perspective of our supersymmetric particle physics based model of inflation? Perhaps surprisingly the answer is positive: we shall show that we can deduce the present day value of dark energy density from the model, together with the measured CMB temperature and the Hubble constant, even if the model does not yet specify the physics of dark energy [1]!

A key point of our approach is that a supersymmetric particle physics based model of inflation enables us to calculate (in principle at least) the (energy or number) densities of all forms of radiation and matter (but excluding dark energy) at some early time t_{RH} after inflation and reheating has taken place, corresponding to the start of the standard hot big bang. For simplicity we consider only one type of matter energy density $\rho_{matter}(t_{RH})$ (which may readily be obtained from the calculated number density) and radiation energy density $\rho_\gamma(t_{RH})$. The argument may be straightforwardly generalised to the case of several components of radiation and matter. Now, using the equations of the standard hot big bang, we wish to obtain their values at the present time t_0 , $\rho_\gamma(t_0)$ and $\rho_{matter}(t_0)$. Without further information this is impossible since we need something to tell us when the present time t_0 is, and moreover the model does not specify either $\rho_{DE}(t_{RH})$, or its equation of state, both of which will influence the evolution of the universe. Therefore let us input into our analysis the present day observed CMB temperature $T_0 = 2.725\text{K}$, which corresponds to a photon density $\rho_\gamma(t_0) = (2.115 \times 10^{-4}\text{eV})^4$, a photon number density $n_\gamma = 410\text{cm}^{-3}$, and, assuming three families of light neutrinos, an entropy density $s = 7.04n_\gamma$. Then, ignoring additional sources of entropy between t_{RH} and t_0 (such as electron-positron annihilation), since we know the equations of state for photons and matter, $\rho_\gamma \sim R^{-4}$ and $\rho_{matter} \sim R^{-3}$, where R is the scale factor of the universe, using the initial values of $\rho_\gamma(t_{RH})$ and $\rho_{matter}(t_{RH})$ predicted by the model and the present value of $\rho_\gamma(t_0)$ from observation, we find $\rho_{matter}(t_0) \approx \rho_{matter}(t_{RH})[\rho_\gamma(t_0)/\rho_\gamma(t_{RH})]^{3/4}$. We emphasise that this determination of $\rho_{matter}(t_0)$ is independent of the unknown dark energy.

Once we have obtained $\rho_{matter}(t_0)$, from a combination of our model calculation and the observed CMB temperature, as outlined above, we now use the observed Hubble constant H_0 , or equivalently the present day critical density ρ_{crit} , to convert $\rho_{matter}(t_0)$ into the various $\Omega_{matter} = \rho_{matter}(t_0)/\rho_{crit}$. Once Ω_{matter} is predicted within some supersymmetric particle physics based model of inflation, supplemented by measurements of the CMB temperature and the Hubble constant, then it is clear that Ω_{DE} is also *predicted* to be

$$\Omega_{DE} = 1 - \Omega_{matter} \quad (5)$$

Thus a model of inflation that is capable of predicting Ω_{matter} using measurements of the CMB temperature and the Hubble constant, is also capable of predicting Ω_{DE} from Eq.5. This sum rule has been written down before, including a curvature term and it has been discussed how to determine each of the components Ω_{DE} and Ω_{matter} from observation.

What we are saying here is quite different from the empirical approach to determining the components of this equation. To begin with we are assuming inflation, so that the curvature contribution is zero. Secondly we are only taking two inputs from observation, namely the CMB temperature and the value of the Hubble constant. Given these inputs we have shown how an inflation model allows us to then *calculate* Ω_{matter} , and hence *deduce* Ω_{DE} from Eq.5.

5 Conclusion

Over the next few years there will be considerable progress in cosmology from the Map and Planck explorer satellites, and in SUSY from the Tevatron and LHC. We believe the time is ripe for a new closer synthesis of SUSY and inflation, and that the most promising scenario will involve these theories meeting at the intermediate scale. We have shown that in this case one may hope to relate different phenomena in cosmology and in particle physics in a much closer and more predictive way than ever before. Finally we have made the original observation that, given the value of the CMB temperature and Hubble constant from observation, an intermediate scale supersymmetric inflation model allows the present day matter relic density to be calculated, and hence the present day DE relic density to be determined from Eq.5 even in the absence of any theory of DE.

ACKNOWLEDGMENTS

I would like to thank co-authors Mar Bastero-Gil, Gordy Kane and David Rayner. I enjoyed conversations at COSMO-01 with many people especially J. Cline, E. Copeland, J. Garcia-Bellido, J.E. Kim, A. Kusenko, A. Liddle, A. Linde, C. Lineweaver, A. Mazumdar, C. Munoz, P. Nilles, K. Olive, J. Peacock, M. Pluemacher and S. Sarkar.

References

- [1] G. L. Kane and S. F. King, arXiv:hep-ph/0109118.
- [2] S. F. King and D. A. Rayner, arXiv:hep-ph/0110349.